

PTPN22 Is Genetically Associated with Risk of Generalized Vitiligo, but CTLA4 Is Not

Greggory S. LaBerge¹, Dorothy C. Bennett², Pamela R. Fain^{1,3} and Richard A. Spritz^{1,4}

Generalized vitiligo is an acquired, multifactorial, polygenic disease in which depigmented spots of skin, overlying hair, and mucous membranes result from autoimmune-mediated loss of melanocytes from affected areas. We examined single-nucleotide polymorphisms (SNPs) in the *PTPN22* and *CTLA4* genes in 126 Caucasian families with multiple cases of generalized vitiligo and associated autoimmune diseases, using a family-based association study design. The *PTPN22* 1858T allele of SNP rs2476601 is significantly associated both with generalized vitiligo and with an expanded autoimmunity phenotype. Individuals carrying the *PTPN22* 1858T allele had an allelic odds ratio (OR) of 2.16 for generalized vitiligo and a genotypic OR of 2.35 as C/T heterozygotes. Similarly, individuals carrying the *PTPN22* 1858T allele had an allelic OR of 2.05 for the expanded autoimmunity phenotype, and a genotypic OR of 2.19 for C/T heterozygotes. Examination of five SNPs in the *CTLA4* gene (rs1863800, rs231775, rs3087243, rs11571302, rs11571297, rs10932037) in the same 126 families yielded no evidence of allelic or genotypic association with either generalized vitiligo or the expanded autoimmune phenotype. These results implicate *PTPN22* in mediating susceptibility to generalized vitiligo and associated autoimmune diseases, but do not support a role for *CTLA4*.

Journal of Investigative Dermatology (2008) **128**, 1757–1762; doi:10.1038/sj.jid.5701233; published online 17 January 2008

INTRODUCTION

Generalized vitiligo is an acquired, non-contagious disorder in which progressive, patchy loss of pigmentation of skin, overlying hair, and mucous membranes results from loss of melanocytes from the involved areas (reviewed in Hann and Nordlund, 2000; Nordlund *et al.*, 2006). Known for thousands of years because of its visually evident phenotype, vitiligo is the most common pigmentation disorder, affecting about 0.4% of Caucasians (Howitz *et al.*, 1977) and occurring with generally similar frequency in other populations (for example, Das *et al.*, 1985a; Lu *et al.*, 2007). Several different etiologic hypotheses have been suggested for generalized vitiligo (reviewed in Nordlund *et al.*, 2006), the most compelling of which involves a combination of unknown environmental and genetic factors interacting to contribute to autoimmune melanocyte destruction.

Most cases of generalized vitiligo occur sporadically, although about 15–20% of patients report one or more affected first-degree relatives. Typically, familial aggregation of generalized vitiligo cases occurs in a non-Mendelian pattern that is suggestive of polygenic, multifactorial inheritance (Mehta *et al.*, 1973; Carnevale *et al.*, 1980; Hafez *et al.*, 1983; Das *et al.*, 1985a,b; Majumder *et al.*, 1988, 1993; Bhatia *et al.*, 1992; Nath *et al.*, 1994; Alkhateeb *et al.*, 2003; LaBerge *et al.*, 2005; Sun *et al.*, 2006), and formal genetic segregation analyses have indicated that multiple major loci contribute to vitiligo susceptibility in a complex interactive manner (Majumder *et al.*, 1993; Nath *et al.*, 1994; Sun *et al.*, 2006). Furthermore, patients with generalized vitiligo are also at increased risk of developing other autoimmune diseases, particularly autoimmune thyroid disease (Graves' disease and autoimmune hypothyroidism), rheumatoid arthritis, psoriasis, latent autoimmune diabetes of adults, pernicious anemia, Addison's disease, and systemic lupus erythematosus (Alkhateeb *et al.*, 2003; LaBerge *et al.*, 2005). These same diseases also occur at increased frequencies in vitiligo patients' first-degree relatives, suggesting that susceptibility to this group of autoimmune diseases is genetically determined.

Genetic linkage and association studies have implicated a number of different genes in susceptibility to generalized vitiligo (reviewed in Spritz, 2007), including several that are thought to play primary roles in the development of autoimmunity. These include the major histocompatibility complex (for example, Foley *et al.*, 1983; Finco *et al.*, 1991; Orecchia *et al.*, 1992; Ando *et al.*, 1993; Schallreuter *et al.*, 1993; al-Fouzan *et al.*, 1995; Zamani *et al.*, 2001; Arcos-

¹Human Medical Genetics Program, University of Colorado Denver, Anschutz Medical Campus, Aurora, Colorado, USA; ²Division of Basic Medical Sciences, St George's, University of London, London, UK; ³Barbara Davis Center for Childhood Diabetes, University of Colorado Denver, Anschutz Medical Campus, Aurora, Colorado, USA and ⁴Department of Pediatrics, University of Colorado Denver, Anschutz Medical Campus, Aurora, Colorado, USA

Correspondence: Professor Richard A. Spritz, Human Medical Genetics Program, University of Colorado Denver, Anschutz Medical Campus, PO Box 6511, MS 8300, Aurora, Colorado 80045 USA.
E-mail: richard.spritz@uchsc.edu

Abbreviations: CI, confidence interval; LD, linkage disequilibrium; OR, odds ratio; SNP, single-nucleotide polymorphism

Received 15 August 2007; revised 20 November 2007; accepted 26 November 2007; published online 17 January 2008

Burgos *et al.*, 2002; Tastan *et al.*, 2004; Fain *et al.*, 2006; Xia *et al.*, 2006; Liu *et al.*, 2007), *CTLA4* (Kemp *et al.*, 1999; Blomhoff *et al.*, 2005; Itirli *et al.*, 2005), and *PTPN22* (Canton *et al.*, 2005), and *NALP1* (Jin *et al.*, 2007a,b). Association of vitiligo with *PTPN22* and *CTLA4* is based on small case-control studies, a study design that is notoriously subject to both false-positive and false-negative errors due to population admixture and stratification (Hirschhorn *et al.*, 2002). To more rigorously test the association of vitiligo with variation in *PTPN22* and *CTLA4*, we carried out family-based association analyses of *CTLA4* and *PTPN22* single-nucleotide polymorphisms (SNPs) in 126 Caucasian families with multiple cases of generalized vitiligo, which included the same families in which we previously showed association to HLA markers (Fain *et al.*, 2006) and *NALP1* (Jin *et al.*, 2007a).

RESULTS

Family-based association analysis of *PTPN22*

We genotyped rs2476601 (*PTPN22* risk allele 1858T) in 712 individuals from 126 Caucasian extended families with multiple cases of vitiligo as well as other autoimmune diseases, derived from the United States and the United Kingdom (Alkhateeb *et al.*, 2003; LaBerge *et al.*, 2005; Jin *et al.*, 2007a). SNP rs2476601 was found to be in Hardy-Weinberg equilibrium in founders (individuals with no parents specified in the pedigree) in the 126 study families (Table 1).

As shown in Table 2, we found that the high-risk 1858T allele of rs2476601 was significantly associated ($P=0.0048$) with generalized vitiligo in the 126 study families. Genotypic analysis showed that the rs2476601 heterozygous C/T genotype was highly associated with vitiligo ($P=0.0012$). The *PTPN22* high-risk 1858T allele showed allelic association with generalized vitiligo [$P=0.024$, odds ratio (OR) 2.16 (95% confidence interval (CI) 1.22–3.82)] (Table 2) from conditional logistic regression analysis.

We also tested an expanded autoimmunity phenotype that included all of the autoimmune diseases with which vitiligo is epidemiologically associated (autoimmune thyroid disease, rheumatoid arthritis, psoriasis, adult-onset autoimmune diabetes mellitus, pernicious anemia, systemic lupus erythematosus, Addison's disease), considering any individual with either vitiligo or any other of these autoimmune diseases as "affected". As shown in Table 2, the rs2476601 1858T allele also showed allelic association ($P=0.0042$) with this expanded autoimmunity phenotype, which was confirmed by conditional logistic regression analysis [$P=0.03$, OR 2.05 (95% CI 1.19–3.53)]. Genotypic analysis, presented in Table 3, showed that the rs2476601 C/T genotype is significantly associated with the expanded autoimmunity phenotype ($P=0.003$), although association with the T/T genotype was not significant ($P=0.750$). Conditional logistic regression analysis provided very similar results, showing significant association of the C/T heterozygote with the autoimmune disease phenotype [$P=0.03$, OR 2.19 (95% CI 1.20–3.97)], but non-significant association of the T/T homozygote [$P=0.14$, OR 3.22 (95% CI 0.68–15.27)].

Table 1. Results of Hardy-Weinberg equilibrium (HWE) analysis of SNP genotypes among founders from 126 multiplex vitiligo-autoimmune disease families

SNP	Observed genotypes in founders	HWE P-value
<i>PTPN22</i>		
rs2476601	T/T 2 C/T 34 C/C 104	0.9170
<i>CTLA4</i>		
rs1863800	T/T 19 C/T 74 C/C 45	0.4210
rs231775	G/G 41 A/G 73 A/A 25	0.7492
rs3087243	A/A 23 A/G 76 G/G 40	0.4304
rs11571302	A/A 24 A/C 75 C/C 38	0.4602
rs11571297	G/G 29 A/G 75 A/A 34	0.5830
rs10932037	T/T 4 C/T 27 C/C 109	0.3750

SNP, single-nucleotide polymorphism.

Family-based association analysis of *CTLA4*

We also examined six SNPs in the *CTLA4* gene (rs1863800, rs231775, rs3087243, rs11571302, rs11571297, rs10932037) for association with generalized vitiligo and the expanded vitiligo-associated autoimmune phenotype in these families. None of the *CTLA4* SNPs tested exhibited either allelic (Table 2) or genotypic (Table 3) association with either generalized vitiligo or with the expanded autoimmunity phenotype. All but one of these SNPs clustered into a single linkage disequilibrium (LD) block (data not shown), and no multiple testing corrections were applied to the nominal P -value threshold ($P=0.05$) due to lack of any apparent association with disease.

DISCUSSION

We have carried out family-based association analyses of *PTPN22* and *CTLA4* SNPs in 126 Caucasian extended families with multiplex cases of vitiligo as well as other autoimmune diseases, confirming the association of generalized vitiligo with the functional *PTPN22* 1858T (R620W)-variant allele (OR 2.16, 95% CI 1.22–3.82) and the heterozygous C/T genotype (OR 2.35, 95% CI 1.25–4.43). Furthermore, we find that the *PTPN22* 1858T allele and C/T genotype are also associated with an expanded autoimmunity phenotype (allelic OR 2.05, 95% CI 1.19–3.53; genotypic OR 2.19, 95% CI 1.20–3.97) that includes generalized vitiligo or

Table 2. Allelic association analyses of *PTPN22* and *CTLA4* SNPs

SNP	FBAT ¹ (P-value)	Regression on alleles (P-value)	Odds ratio (95% CI)
<i>Vitiligo</i>			
<i>PTPN22</i>			
rs2476601	0.0048 ²	0.024 ²	2.16 (1.22–3.82)
<i>CTLA4</i>			
rs1863800	0.3742	0.495	0.89 (0.63–1.24)
rs231775	0.5440	0.592	1.09 (0.79–1.52)
rs3087243	0.3518	0.683	0.934 (0.67–1.30)
rs11571302	0.3628	0.930	0.986 (0.71–1.36)
rs11571297	0.3942	0.858	1.03 (0.74–1.44)
rs10932037	0.8688	1.00	1.00 (0.61–1.65)
<i>Expanded autoimmunity phenotype</i>			
<i>PTPN22</i>			
rs2476601	0.0042 ²	0.03 ²	2.05 (1.19–3.53)
<i>CTLA4</i>			
rs1863800	0.2907	0.305	0.85 (0.62–1.16)
rs231775	0.4198	0.310	0.85 (0.62–1.16)
rs3087243	0.2460	0.370	0.87 (0.63–1.18)
rs11571302	0.2655	0.578	0.92 (0.67–1.25)
rs11571297	0.2920	0.465	1.13 (0.82–1.56)
rs10932037	0.8858	0.809	0.94 (0.59–1.52)

CI, confidence interval; FBAT, family-based association test; SNP, single-nucleotide polymorphism.

¹P-values based on FBAT additive model.

²P-values corrected for three independent tests: two LD blocks in *CTLA4* and one SNP in *PTPN22*.

any of the other autoimmune diseases with which vitiligo is epidemiologically associated (Alkhateeb *et al.*, 2003; LaBerge *et al.*, 2005). These findings strongly support a causal role for the *PTPN22* 1858T allele of rs2476601 (or another variant with which 1858T is in close LD) in susceptibility to both generalized vitiligo and these other autoimmune diseases. We found no association of the *PTPN22* rs2476601 homozygous T/T genotype with either generalized vitiligo or the expanded autoimmune phenotype, most likely due to limited power resulting from the low frequency of the T allele (0.13 in this study) and the consequent small number of T/T homozygotes observed (8 of 712 individuals genotyped).

However, our analyses of the *CTLA4* SNPs showed no apparent allelic or genotypic association with either generalized vitiligo or the expanded autoimmunity phenotype. These results are in contrast with previous reports of genetic association of *CTLA4* markers with vitiligo based on small case-control studies (Kemp *et al.*, 1999; Blomhoff *et al.*, 2005; Itirli *et al.*, 2005). Kemp *et al.* (1999) reported the association of the 106-bp allele of a microsatellite polymorphism in the *CTLA4* 3'-untranslated region with vitiligo (especially autoimmune-associated) in 74 European Caucasian patients *versus* 173 controls. The same group (Blomhoff *et al.*, 2005) subsequently reported the association

of *CTLA4* SNPs MH30 (23 kb 5' of the *CTLA4* gene), rs3087243, rs11571302, and rs7565213 with vitiligo in 27 European Caucasian patients with associated autoimmune disease *versus* 140 controls; it is not clear whether the case and control cohorts in these two studies were independent. Itirli *et al.* (2005) reported the association of the 96-bp allele of the same *CTLA4* microsatellite in 36 Turkish patients *versus* 100 controls. It may be that these previous "associations" represent spurious false-positive results, due to either occult population stratification (reviewed in Hirschhorn *et al.*, 2002) or the very small sample size and highly unbalanced numbers of cases and controls in both studies. Furthermore, Blomhoff *et al.* (2005) did not correct for multiple testing, which would have rendered the reported *CTLA4* associations insignificant. Itirli *et al.* (2005) reported significance based on the observation of only four vitiligo patients *versus* two controls with the 96-bp allele, whereas the 112-bp allele, which showed the largest difference in allele frequency between cases and controls, was not significant, indicating that "significance" of the 96-bp allele almost certainly represents statistical fluctuation due to the very small number of observations.

PTPN22 and *CTLA4* are thought to function as general autoimmunity susceptibility loci (Brand *et al.*, 2005; Gregersen *et al.*, 2006). The *PTPN22* 1858T variant results in an arginine to tryptophan substitution that disrupts interaction between Lyp and Csk protein tyrosine kinases, dis-inhibiting T-cell activation (Siminovitch, 2004) and perhaps thereby increasing susceptibility to autoimmune disease (Brand *et al.*, 2005). The *PTPN22* 1858T variant has been associated with type 1 diabetes, rheumatoid arthritis, systemic lupus erythematosus, Graves' disease (Bottini *et al.*, 2004; Siminovitch, 2004; Criswell *et al.*, 2005; Zhernakova *et al.*, 2005), and vitiligo (Canton *et al.*, 2005). In contrast, studies of patients with psoriasis and multiple sclerosis showed no association with the *PTPN22* risk allele in two family-based studies (Criswell *et al.*, 2005; Nistor *et al.*, 2005), suggesting that *PTPN22* may not truly be associated with these diseases.

Our study utilized a family-based design to test association of *PTPN22* and *CTLA4* with generalized vitiligo, an approach that is more rigorous than previous case-control studies of these genes in vitiligo. Our findings demonstrate that *PTPN22* can be included along with HLA and *NALP1* among those genes confirmed to play a role in polygenic susceptibility to generalized vitiligo, but do not support a role for *CTLA4*.

MATERIALS AND METHODS

Subjects

Peripheral blood or saliva samples were obtained from 712 individuals from 126 extended families with multiplex cases of generalized vitiligo as well as other autoimmune diseases, derived from the United States and United Kingdom. All families were of Caucasian origin (as self-reported) and had two or more family members with generalized vitiligo and at least one family member with one or more other autoimmune disease with which generalized vitiligo is epidemiologically associated (autoimmune thyroid disease, latent autoimmune diabetes in adults, psoriasis, pernicious anemia, systemic lupus erythematosus, rheumatoid arthritis,

Table 3. Genotypic association analyses of *PTPN22* and *CTLA4* SNPs

SNP	Genotype	FBAT ¹ (P-value)	Regression on genotypes (P-value)	Odds ratio (95% CI)
<i>Vitiligo</i>				
<i>PTPN22</i>				
rs2476601	T/T	0.814	0.126	3.42 (0.71–16.54)
	C/T	0.0012 ²	0.024 ²	2.35 (1.25–4.43)
	C/C	0.999	Referent	
<i>CTLA4</i>				
rs1863800	T/T	0.807	0.542	0.797 (0.385–1.65)
	C/T	0.488	0.576	0.886 (0.579–1.35)
	C/C	0.327	Referent	
rs231775	A/A	0.408	0.592	0.830 (0.419–1.64)
	A/G	0.563	0.762	1.08 (0.657–1.77)
	G/G	0.944	Referent	
rs3087243	A/A	0.732	0.652	0.854 (0.430–1.70)
	A/G	0.552	0.922	0.978 (0.623–1.53)
	G/G	0.309	Referent	
rs11571302	A/A	0.448	0.752	0.892 (0.440–1.81)
	A/C	0.989	0.523	1.16 (0.738–1.82)
	C/C	0.528	Referent	
rs11571297	G/G	0.567	0.706	1.15 (0.553–2.40)
	A/G	0.957	0.325	1.36 (0.740–2.48)
	A/A	0.473	Referent	
rs10932037	T/T	0.848	1.00	1.00 (0.234–4.28)
	C/T	0.751	1.00	1.00 (0.572–1.75)
	C/C	0.799	Referent	
<i>Expanded autoimmunity phenotype</i>				
<i>PTPN22</i>				
rs2476601	T/T	0.750	0.140	3.22 (0.68–15.27)
	C/T	0.003 ²	0.03 ²	2.19 (1.20–3.97)
	C/C	0.999	Referent	
<i>CTLA4</i>				
rs1863800	T/T	0.636	0.379	0.734 (0.368–1.46)
	C/T	0.553	0.358	0.829 (0.555–1.24)
	C/C	0.299	Referent	
rs231775	A/A	0.187	0.310	0.712 (0.369–1.37)
	A/G	0.271	0.933	1.02 (0.638–1.63)
	G/G	0.896	Referent	
rs3087243	A/A	0.557	0.372	0.748 (0.394–1.42)
	A/G	0.619	0.550	0.876 (0.568–1.35)
	G/G	0.256	Referent	
rs11571302	A/A	0.371	0.506	0.801 (0.416–1.54)
	A/C	0.992	0.955	1.01 (0.656–1.56)
	C/C	0.426	Referent	
rs11571297	G/G	0.454	0.418	1.32 (0.672–2.60)
	A/G	0.987	0.334	1.31 (0.759–2.24)
	A/A	0.403	Referent	
rs10932037	T/T	0.784	0.942	0.948 (0.223–4.02)
	C/T	0.965	0.789	0.930 (0.549–1.58)
	C/C	0.955	Referent	

CI, confidence interval; FBAT, family-based association test; SNP, single-nucleotide polymorphism.

¹P-values based on FBAT genotype model.

²P-values corrected for 3 independent tests: 2 LD blocks in *CTLA4* and 1 SNP in *PTPN22*.

Addison's disease; Alkhateeb *et al.*, 2003; LaBerge *et al.*, 2005). Diagnostic criteria for generalized vitiligo were consistent with those of the Vitiligo European Task Force (Ta'ieb and Picardo, 2007). Exclusion criteria were atypical lesion distribution, congenital or static skin depigmentation (for example, birthmarks, piebaldism, Waardenburg syndrome), depigmentation secondary to the use of melanocytotoxic chemicals, inflammatory skin diseases (systemic lupus erythematosus, lichen planus, psoriasis), and post-infectious or post-traumatic causes.

All available affected and unaffected family members completed a clinical history questionnaire reporting vitiligo age of onset and course of treatment, and a checklist of approximately 50 autoimmune and autoinflammatory diseases. Each vitiligo patient completed a skin-lesion map. All data were reviewed by study investigators and staff, and individuals in whom diagnoses were uncertain based on standard diagnostic criteria (Nordlund *et al.*, 2006) were excluded from the study. This study conformed to the Declaration of Helsinki Principles and was approved by the Colorado Multiple Institutional Review Board and the South East Research Ethics Committee. Written, informed consent was provided by all study participants.

SNP genotyping

DNA was isolated from peripheral blood using a genomic DNA purification kit (Purgene, Gentra Systems) or from saliva obtained using the DNA self-collection kit (Oragene; DNA Genotek, Ottawa, Ontario, Canada). Genotyping was carried out for *PTPN22*-rs2476601 (1858C/T), and *CTLA4*-rs1863800, rs231775 (+49G), rs3087243 (CT60), rs11571302 (JO31), rs11571297 (JO27_1), rs10932037 (CTIC154_1) in 10-μl PCR reaction volumes with 20 ng genomic DNA using standard methods. SNP alleles were detected with the Applied Biosystems (ABI, Foster City, CA) PRISM SNaPshot Multiplex Kit using an ABI 3130 Genetic Analyzer. Automated genotyping was performed using ABI Genemapper version 3.7 software; all genotypes were manually checked for accuracy and allele calling consistency. Mendelian inheritance of all markers was checked through all levels of PedCheck (O'Connell and Weeks, 1998). Haplotype analysis of *CTLA4* SNPs was carried out using the error-screening routine of Merlin, version 1.0.0 (Abecasis *et al.*, 2002).

Statistical analyses

Deviations from expected Hardy-Weinberg proportions were tested for each SNP in founders (no parents in the pedigree) and in persons not in the lineage in all 126 families using χ^2 -analysis, considering $P < 0.05$ significant. Calculation of LD between *CTLA4* SNPs was performed using Haploview software (Barrett *et al.*, 2005), version 3.32, with haplotype blocks being determined by the method of Gabriel *et al.* (2002). Single-locus association analyses of each SNP were carried out using the family-based association test (Horvath *et al.*, 2001). Conditional logistic regression analysis was carried out using STATA, version 9.2, which uses family-based data to create a matched pseudo case-control data set, test for allelic or genotypic association and estimate an OR for disease risk (Cordell and Clayton, 2002). Nominally significant P -values (< 0.05) were corrected for three independent tests based on LD analysis, using Haploview (two LD blocks defined in the *CTLA4* gene and one SNP in *PTPN22*).

CONFLICT OF INTEREST

The authors state no conflict of interest.

ACKNOWLEDGMENTS

We thank the many families who participated in this study, and the UK Vitiligo Society, the US National Vitiligo Foundation, and Vitiligo Support International for their enthusiastic help in family ascertainment. We thank Anita Amadi-Myers and Paulene Holland for their invaluable assistance. This work was supported by grants AR45584, AI46374, and DK57538 from the National Institutes of Health.

REFERENCES

- Abecasis GR, Cherny SS, Cookson WO, Cardon LR (2002) Merlin-rapid analysis of dense genetic maps using sparse gene flow trees. *Nat Genet* 30:97–101
- al-Fouzan A, al-Arbash M, Fouad F, Kaaba SA, Mousa MA, al-Harbi SA (1995) Study of HLA class I/II and T lymphocyte subsets in Kuwaiti vitiligo patients. *Eur J Immunogenet* 22:209–13
- Alkhateeb A, Fain PR, Thody A, Bennett DC, Spritz RA (2003) Epidemiology of vitiligo and associated autoimmune diseases in Caucasian probands and their families. *Pigment Cell Res* 16:208–14
- Ando I, Chi HI, Nakagawa H, Otsuka F (1993) Difference in clinical features and HLA antigens between familial and non-familial vitiligo of non-segmental type. *Br J Dermatol* 129:408–10
- Arcos-Burgos M, Parodi E, Salgar M, Bedoya E, Builes J, Jaramillo D *et al.* (2002) Vitiligo: complex segregation and linkage disequilibrium analyses with respect to microsatellite loci spanning the HLA. *Hum Genet* 110:334–42
- Barrett JC, Fry B, Maller J, Daly MJ (2005) Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics* 21:263–5
- Bhatia PS, Mohan L, Pandey ON, Singh KK, Arora SK, Mukhija RD (1992) Genetic nature of vitiligo. *J Dermatol Sci* 4:180–4
- Blomhoff A, Kemp EH, Gawkrödger DJ, Weetman AP, Husebye ES, Akselsen HE *et al.* (2005) *CTLA4* polymorphisms are associated with vitiligo, in patients with concomitant autoimmune diseases. *Pigment Cell Res* 18:55–8
- Bottini N, Musumeci L, Alonso A, Rahmouni S, Nika K, Rostamkhani M *et al.* (2004) A functional variant of lymphoid tyrosine phosphatase is associated with type I diabetes. *Nat Genet* 36:337–8
- Brand O, Gough S, Heward J (2005) HLA, *CTLA-4* and *PTPN22*: the shared genetic master-key to autoimmunity? *Expert Rev Mol Med* 7:1–15
- Canton I, Askar S, Gavalas NG, Gawkrödger DJ, Blomhoff A, Watson PF *et al.* (2005) A single-nucleotide polymorphism in the gene encoding lymphoid protein tyrosine phosphatase (*PTPN22*) confers susceptibility to generalised vitiligo. *Genes Immun* 6:584–7
- Carnevale A, Zavala C, del Castillo V, Ruiz Maldonado R, Tamayo L (1980) Genetic analysis of 127 families with vitiligo. *Rev Invest Clin* 32:37–41
- Cordell HJ, Clayton DG (2002) A unified stepwise regression procedure for evaluating the relative effects of polymorphisms within a gene using case/control or family data: application to HLA in type 1 diabetes. *Am J Hum Genet* 70:124–41
- Criswell LA, Pfeiffer KA, Lum RF, Gonzales B, Novitzke J, Kern M *et al.* (2005) Analysis of families in the multiple autoimmune disease genetics consortium (MADGC) collection: the *PTPN22* 620W allele associates with multiple autoimmune phenotypes. *Am J Hum Genet* 76:561–71
- Das SK, Majumder PP, Chakraborty R, Majumdar TK, Haldar B (1985a) Studies on vitiligo I. Epidemiological profile in Calcutta, India. *Genet Epidemiol* 2:71–8
- Das SK, Majumder PP, Majumdar TK, Haldar B (1985b) Studies on vitiligo II. Familial aggregation and genetics. *Genet Epidemiol* 2:255–62
- Fain PR, Babu SR, Bennett DC, Spritz RA (2006) HLA class II haplotype DRB1*04-DQB1*0301 contributes to risk of familial generalized vitiligo and early disease onset. *Pigment Cell Res* 19:51–7
- Finco O, Cuccia M, Martinetti M, Ruberto G, Orecchia G, Rabbiosi G (1991) Age of onset in vitiligo: relationship with HLA supratypes. *Clin Genet* 39:48–54
- Foley LM, Lowe NJ, Misheloff E, Tiwari JL (1983) Association of HLA-DR4 with vitiligo. *J Am Acad Dermatol* 8:39–40
- Gabriel SB, Schaffner SF, Nguyen H, Moore JM, Roy J, Blumenstiel B *et al.* (2002) The structure of haplotype blocks in the human genome. *Science* 296:2225–9
- Gregersen PK, Lee H-S, Batliwalla F, Begovich AB (2006) *PTPN22*: setting thresholds for autoimmunity. *Semin Immunol* 18:214–23
- Hafez M, Sharaf L, Abd el-Nabi SM (1983) The genetics of vitiligo. *Acta Derm Venereol* 63:249–51
- Hann SK, Nordlund JJ (2000) *Vitiligo: a Comprehensive Monograph on Basic and Clinical Science*. Oxford: Blackwell Science
- Hirschhorn JN, Lohmueller K, Byrne E, Hirschhorn K (2002) A comprehensive review of genetic association studies. *Genet Med* 4:45–61
- Horvath S, Wei E, Xu X, Palmer LJ, Baur M (2001) Family-based association test method: age of onset traits and covariates. *Genet Epidemiol* 21(Suppl 1): S403–8
- Howitz J, Brodthagen H, Schwartz M, Thomsen K (1977) Prevalence of vitiligo. Epidemiological survey on the Isle of Bornholm, Denmark. *Arch Dermatol* 113:47–52
- Itirli G, Pehlivan M, Alper S, Yuksel SE, Onay H, Ozkinay F *et al.* (2005) Exon-3 polymorphism of *CTLA-4* gene in Turkish patients with vitiligo. *J Dermatol Sci* 38:225–7
- Jin Y, Birlea SA, Fain PR, Spritz RA (2007b) Genetic variations in *NALP1* are associated with generalized vitiligo in a Romanian population. *J Invest Dermatol* 127:2558–62
- Jin Y, Mailloux CM, Gowan K, Riccardi SL, LaBerge G, Bennett DC *et al.* (2007a) *NALP1* in vitiligo-associated multiple autoimmune disease. *N Engl J Med* 356:1216–25
- Kemp EH, Ajjan RA, Waterman EA, Gawkrödger DJ, Cork MJ, Watson PF *et al.* (1999) Analysis of a microsatellite polymorphism of the cytotoxic T-lymphocyte antigen-4 gene in patients with Vitiligo. *Br J Dermatol* 140:73–8
- LaBerge G, Mailloux CM, Gowan K, Holland P, Bennett DC, Fain PR *et al.* (2005) Early disease onset and increased risk of other autoimmune diseases in familial generalized vitiligo. *Pigment Cell Res* 18:300–5
- Liu JB, Li M, Chen H, Zhong SQ, Yang S, Du WD *et al.* (2007) Association of vitiligo with HLA-A2: a meta-analysis. *J Eur Acad Dermatol Venereol* 21:205–13
- Lu T, Gao T, Wang A, Jin Y, Li Q, Li C (2007) Vitiligo prevalence study in Shaanxi Province, China. *Int J Dermatol* 46:47–51
- Majumder PP, Das SK, Li CC (1988) A genetical model for vitiligo. *Am J Hum Genet* 43:119–25
- Majumder PP, Nordlund JJ, Nath SK (1993) Pattern of familial aggregation of vitiligo. *Arch Dermatol* 129:994–8
- Mehta NR, Shah KC, Theodore C, Vyas VP, Patel AB (1973) Epidemiological study of vitiligo in Surat area, South Gujarat. *Indian J Med Res* 61: 145–54
- Nath SK, Majumder PP, Nordlund JJ (1994) Genetic epidemiology of vitiligo: multilocus recessivity cross-validated. *Am J Hum Genet* 55:981–90
- Nistor I, Nair RP, Stuart P, Hiremagalore R, Thompson RA, Jenisch S *et al.* (2005) Protein tyrosine phosphatase gene *PTPN22* polymorphism in psoriasis: lack of evidence for association. *J Invest Dermatol* 125:395–6
- Nordlund JJ, Ortonne J-P, LePoole IC (2006) Vitiligo vulgaris. In: *The Pigmentary System* (Nordlund JJ, Boissy RE, Hearing VJ, King, RA, Oetting, WS, Ortonne J-P, eds), 2nd ed. Malden, MA: Blackwell Publishing, 551–98
- O'Connell JR, Weeks DE (1998) PedCheck: a program for identification of genotype incompatibilities in linkage analysis. *Am J Hum Genet* 63:259–66
- Orecchia G, Perfetti L, Malagoli P, Borghini F, Kipervarg Y (1992) Vitiligo is associated with a significant increase in HLA-A30, Cw6 and DQw3 and a decrease in C4AQ0 in northern Italian patients. *Dermatology* 185:123–7

- Schallreuter KU, Levenig C, Kuhl P, Loliger C, Hohl-Tehari M, Berger J (1993) Histocompatibility antigens in vitiligo: Hamburg study on 102 patients from northern Germany. *Dermatology* 187:186–92
- Siminovitch KA (2004) *PTPN22* and autoimmune disease. *Nat Genet* 36:1248–9
- Spritz RA (2007) The genetics of generalized vitiligo and associated autoimmune disease. *Pigment Cell Res* 20:271–8
- Sun X, Xu A, Wei X, Ouyang J, Lu L, Chen M *et al.* (2006) Genetic epidemiology of vitiligo: a study of 815 probands and their families from south China. *Int J Dermatol* 45:1176–81
- Taïeb A, Picardo M, VETF Members (2007) The definition and assessment of vitiligo: a consensus report of the Vitiligo European Task Force. *Pigment Cell Res* 20:27–35
- Tastan HB, Akar A, Orkunoglu FE, Arca E, Inal A (2004) Association of HLA class I antigens and HLA class II alleles with vitiligo in a Turkish population. *Pigment Cell Res* 17:181–4
- Xia Q, Zhou WM, Liang YH, Ge HS, Liu HS, Wang JY *et al.* (2006) MHC haplotypic association in Chinese Han patients with vitiligo. *J Eur Acad Dermatol Venereol* 20:941–6
- Zamani M, Spaepen M, Sghar SS, Huang C, Westerhof W, Nieuweboer-Krobotova L *et al.* (2001) Linkage and association of HLA class II genes with vitiligo in a Dutch population. *Br J Dermatol* 145:90–4
- Zhernakova A, Eerligh P, Wijmenga C, Barrera P, Roep BO, Koeleman BP (2005) Differential association of the *PTPN22* coding variant with autoimmune diseases in a Dutch population. *Genes Immun* 6:459–61